arXiv:1210.0046v1 [math.CA] 28 Sep 2012

ON FUNCTIONAL INEQUALITIES FOR THE PSI FUNCTION

BARKAT ALI BHAYO

Abstract. The volume of the unit ball in $\mathbb{R}^n$ can be expressed in terms of the gamma function. This connection has lead in the recent years to questions motivated by geometric considerations involving the gamma function and other special functions. Here we give some functional inequalities for the psi function, the logarithmic derivative of the gamma function.

2010 Mathematics Subject Classification: 33B15
Keywords and phrases: Gamma function, psi function, inequalities.

1. Introduction

For $\mathrm{Re}\,x > 0$, $\mathrm{Re}\,y > 0$, we define the classical *gamma function* $\Gamma(x)$ and *psi function* $\psi(x)$ by

$$\Gamma(x) = \int_0^\infty e^{-t}t^{x-1}\,dt,\ \psi(x) = \frac{\Gamma'(x)}{\Gamma(x)},$$

respectively. The recurrence relations of Γ and ψ are

$$\Gamma(1+x) = x\Gamma(x), \quad \psi(x+1) = \frac{1}{x} + \psi(x).$$

Note that

$$\psi(1) = -\gamma \quad \text{and} \quad \psi(1/2) = -2\log 2 - \gamma$$

where γ is the *Euler-Mascheroni* constant. Throughout this paper, we denote by $c = 1.4616\ldots$ the only positive root of the the equation $\psi(x) = 0$ (see [AS, 6.3.19]).

1.1. **Lemma.** [S, Theorem 2.3] *For all $a \geq 1$, the function*

$$f(x) = \frac{\Gamma(1+x)^a}{\Gamma(1-ax)}$$

is decreasing function of $x \geq 0$.

The following result is the counterpart of the Lemma 1.1.

`File: psi120929.tex, printed: 2012-10-2,`

1.2. **Theorem.** *For* $a > 2, b > 1$ *with* $a > b$, *the following function*

$$f(x) = \frac{\psi(1+bx)^a}{\psi(1+ax)^b}$$

is increasing (decreasing) in $(1/2, \infty)$ *for* $a > 2, b > 1, a > b$ $(a > 1, b > 2, a < b)$.

In particular,

(1) $\left(\frac{\psi(1+b/2)}{\psi(1+bx)}\right)^a < \left(\frac{\psi(1+a/2)}{\psi(1+ax)}\right)^b, \quad a > 2,\ b > 1,\ a > b,$

(2) $\left(\frac{\psi(1+bx)}{\psi(1+b/2)}\right)^a < \left(\frac{\psi(1+ax)}{\psi(1+a/2)}\right)^b, \quad a > 1,\ b > 2,\ a < b,$

(3) $C\psi(1+2x) < \psi(1+x)^2$,
where $C = \frac{(2-\log 4-\gamma)^2}{1-\gamma} \approx 0.0031\ldots$.

The functional inequality of Theorem 1.3 is reminiscent of Theorem 5.12 of [AVV].

1.3. **Theorem.** *The function* $f(x) = 1/\psi(\cosh(x))$ *is decreasing and convex from* $(1, \infty)$ *onto* $(0, a)$, *where* $f(1) = 1/\psi(\cosh(1)) = 13.1559\ldots = a$. *In particular,*

$$\frac{2\psi(r)\psi(s)}{\psi(\sqrt{(1+rs+r's')/(rs)})} \leq \psi(r) + \psi(s),$$

for all $r, s \in (1, \infty)$ *and* $z' = \sqrt{1-z^2}$.

The volume of the unit ball $\mathbb{B}^n$ in $\mathbb{R}^n$ is defined as

$$\Omega_n = \frac{\pi^{n/2}}{\Gamma(n/2+1)} = \prod_{j=1}^{n} \int_0^{\pi} (\sin t)^j dt, \quad \Omega_0 = 1, \quad n \in \mathbb{N},$$

(see [AVV, pp. 37]).

1.4. **Theorem.** *The following inequalities hold*

(1) $\sqrt{\Omega_{2n}\Omega_{2(n-1)}} \leq \Omega_{(2n-1)}$,

(2) $\left(\frac{\Omega_{n-1}}{\Omega_n}\right)^k \leq \frac{\Omega_{k(n-1)}}{\Omega_{kn}}$,

for all $k, n \in \mathbb{N}$.

2. PRELIMINARIES AND PROOFS

For the following lemma see [A1, EL, A3, EGP]

2.1. **Lemma.** *For* $x > 0$ *we have*

(1) $\log x - \frac{1}{x} < \psi(x) < \log x - \frac{1}{2x}$,

(2) $\psi'(x) > \dfrac{1}{x} + \dfrac{1}{2x^2}$,

(3) $\psi''(x) < \dfrac{1}{x} - 2\psi'(x)$,

(4) $(\psi'(x))^2 + \psi''(x) > 0$,

(5) $\psi'(x)e^{\psi(x)} < 1$.

2.2**. Proof of Theorem 1.2** Let

$$g(x) = \log f(x) = a\log(\psi(1+bx)) - b\log(\psi(1+ax)).$$

Differentiating with respect to x we get

$$g'(x) = ab\left(\frac{\psi'(1+bx)}{\psi(1+bx)} - \frac{\psi'(1+ax)}{\psi(1+ax)}\right).$$

By Lemma 2.1, ψ' and ψ are decreasing and increasing, respectively. Clearly

$$\psi'(1+bx)\psi(1+ax) - \psi'(1+ax)\psi(1+bx) > 0,$$

hence f is increasing. Letting $a = 2$, $b = 1$ we get

$$f(1/2) < f(x),$$

and this completes the proof. □

2.3. **Lemma.** *The function*

$$f(x) = \frac{\psi'(\cosh(x))\sinh(x)}{\psi(\cosh(x))^2},$$

is decreasing in $x \in (1, \infty)$.

Proof. Letting $r = \cosh(x)$,

$$f(x) = \frac{\sqrt{r^2-1}\,\psi'(r)}{(\psi(r))^2}.$$

We get

$$\begin{aligned} f'(x) &= \frac{-(r^2-1)\psi'(r)^2 + \psi(r)\left(r\psi'(r) + (r^2-1)\,\psi''(r)\right)}{\sqrt{r^2-1}\,\psi(r)^2} \\ &< \frac{(r^2-1)\,\psi''(r) + \psi(r)\left(r\psi'(r) + (r^2-1)\,\psi''(r)\right)}{\sqrt{r^2-1}\,\psi(r)^2} < 0 \end{aligned}$$

by Lemma 2.1(4). □

2.4**. Proof of Theorem 1.3.** Differentiating with respect to x we get

$$f'(x) = -\frac{\psi'(\cosh(x))\sinh(x)}{\psi(\cosh(x))^2},$$

which is negative and decreasing, hence f is convex. This implies that

$$\frac{1}{2}\left(\frac{1}{\psi(\cosh(x))}+\frac{1}{\psi(\cosh(y))}\right)\leq\frac{1}{\psi(\cosh(x+y)/2)}.$$

The inequality follows if we let $r=\cosh(x), s=\cosh(y)$ and use $\cosh^2((x+y)/2)=(1+xy+x'y')/(xy)$. $\square$

For convenience we use the notation $\mathbb{R}_+=(0,\infty)$.

2.5. **Lemma.** [N, Thm 2.1] *Let* $f:\mathbb{R}_+\to\mathbb{R}_+$ *be a differentiable, log-convex function and let* $a\geq 1$*. Then* $g(x)=(f(x))^a/f(a\,x)$ *decreases on its domain. In particular, if* $0\leq x\leq y$ *, then the following inequalities*

$$\frac{(f(y))^a}{f(a\,y)}\leq\frac{(f(x))^a}{f(a\,x)}\leq(f(0))^{a-1}$$

hold true. If $0<a\leq 1$*, then the function* g *is an increasing function on* $\mathbb{R}_+$ *and inequalities are reversed.*

2.6. **Corollary.** *For* $k>1$ *and* $c<x\leq y$*, the following inequality holds*

$$\left(\frac{\psi(x)}{\psi(y)}\right)^k\leq\frac{\psi(kx)}{\psi(ky)}.$$

Proof. Let $g(x)=\log(1/\psi(x))$. Differentiating with respect to x and by Lemma 2.1 we get

$$\begin{aligned}
g''(x) &= \frac{\psi'(x)^2-\psi(x)\psi(x)^2}{\psi(x)^2}\\
&> \frac{\psi'(x)^2-(\log(x)-1/(2x))(1/x-2\psi'(x))}{\psi(x)^2}\\
&> \frac{\psi'(x)^2-(\log(x)-1/(2x))(1/x-2(1/x+1/(2x^2)))}{\psi(x)^2}\\
&> \frac{\psi'(x)^2+(\log(x)-1/(2x))(1/x+1/x^2)}{\psi(x)^2}>0.
\end{aligned}$$

This implies that g is convex. Now the rest of proof follows easily from Lemma 2.5. $\square$

2.7. **Corollary.** *The function* $f(x)=\operatorname{artanh}(\psi(\tanh(x)))$ *is strictly increasing and concave from* (c,∞) *onto* (l,m)*, where*

$$f(c)=\operatorname{artanh}(\psi(\tanh(c)))=-0.9934\ldots=l$$

and

$$f(\infty)=-\operatorname{artanh}(\gamma)=-0.6582\ldots=m.$$

In particular,

(1) $\psi\left(\dfrac{r+s}{1+rs+r's'}\right) > \dfrac{\psi(r)+\psi(s)}{1+\psi(r)\psi(s)+\sqrt{1-\psi(r)^2}\sqrt{1-\psi(s)^2}}$,
for all $r, s \in (0,1)$,

(2) $\psi\left(\dfrac{r+s}{1+rs}\right) > \tanh\left(\dfrac{1}{2}\,\mathrm{artanh}\left(\dfrac{\psi(2r)+\psi(2s)}{1+\psi(2r)\psi(2s)}\right)\right)$,
for all $r, s \in (0,1)$,

(3) $\dfrac{1+\psi(\tanh(r))}{1-\psi(\tanh(r))}\dfrac{1-\psi(\tanh(s))}{1+\psi(\tanh(s))} < e^{2a(r-s)}$,
for all $r, s \in (c,\infty)$, *where*

$$f'(c) = \frac{\psi'(\tanh(c))\mathrm{sech}^2(c)}{1-\psi(\tanh(c))^2} = 0.8807\ldots = a.$$

Proof. Differentiating f with respect to x, we get

$$f'(x) = \frac{\psi'(\tanh(x))\mathrm{sech}^2(x)}{1-\psi(\tanh(x))^2} = \frac{F(x)}{G(x)}.$$

Which is positive and decreasing, because

$$F'(x) = \psi''(\tanh(x))\mathrm{sech}^4(x) - 2\psi'(\tanh(x))\mathrm{sech}^2(x)\tanh(x) < 0.$$

Clearly $G(x)$ is increasing, hence f is concave. The concavity of the function implies that

$$f\left(\frac{x+y}{2}\right) > \frac{f(x)+f(y)}{2}$$

$$\iff \psi\left(\tanh\left(\frac{x+y}{2}\right)\right) > \tanh\left(\frac{\mathrm{artanh}(\psi(\tanh(x)))+\mathrm{artanh}(\psi(\tanh(y)))}{2}\right).$$

We get (1) by using

$$\tanh\left(\frac{x+y}{2}\right) = \frac{\tanh(x+y)}{1+\sqrt{1-\tanh^2(x+y)}} = \frac{r+s}{1+rs+r's'},$$

$$\tanh\left(\frac{u+v}{2}\right) = \frac{R+S}{1+RS+R'S'},$$

and letting $r=\tanh(x)$, $s=\tanh(y)$, $R=\psi(r)$, $S=\psi(s)$, $u=\mathrm{artanh}(\psi(\tanh(x)))$ and $v=\mathrm{artanh}(\psi(\tanh(y)))$. For (2),

$$\begin{aligned}\mathrm{artanh}\psi\left(\tanh\left(\frac{x+y}{2}\right)\right) &> \frac{\mathrm{artanh}(\psi(\tanh(x)))+\mathrm{artanh}(\psi(\tanh(y)))}{2}\\ &= \frac{1}{2}\mathrm{artanh}\left(\frac{\psi(\tanh(x))+\psi(\tanh(y))}{1+\psi(\tanh(x))\psi(\tanh(y))}\right).\end{aligned}$$

Letting $r=\tanh(x/2)$ and $s=\tanh(y/2)$ we get (2). The derivative $f'(x)$ tends to a when x tends to c. By Mean Value Theorem we get $f(r)-f(s)<a(r-s)$. This is equivalent to

$$\frac{1}{2}\log\left(\frac{1+\psi(\tanh(r))}{1-\psi(\tanh(r))}\right)-\frac{1}{2}\log\left(\frac{1+\psi(\tanh(s))}{1-\psi(\tanh(s))}\right)<a(r-s),$$

hence (3) follows, and this completes the proof. ☐

2.8. **Lemma.** [KMSV, Thm 1.7] *Let $f:\mathbb{R}_+\to\mathbb{R}_+$ be a differentiable function and for $c\neq 0$ define*

$$g(x)=\frac{f(x^c)}{(f(x))^c}\,.$$

We have the following

(1) *if $h(x)=\log(f(e^x))$ is a convex function, then $g(x)$ is monotone increasing for $c,x\in(0,1)$ or $c,x\in(1,\infty)$ or $c<0,x>1$and monotone decreasing for $c\in(0,1),x>1$ or $c>1$, $x\in(0,1)$ or $c<0,x\in(0,1)$,*

(2) *if $h(x)$ is a concave function, then $g(x)$ is monotone increasing for $c\in(0,1),x>1$ or $c>1$, $x\in(0,1)$ or $c<0$, $x\in(0,1)$ and monotone decreasing for $c,x\in(0,1)$ or $c>1,x>1$ or $c<0$, $x>1$.*

2.9. **Corollary.** *The following inequalities hold for $r,s\in(c,\infty)$,*

(1) $\psi(\sqrt{rs})\geq\sqrt{\psi(r)\psi(s)}$,
equality holds with $r=s$,
(2) $\psi(x^k)<\psi(r)^k, \quad k\in(0,1)$,
(3) $\psi(r)^k<\psi(r^k), \quad k>1$.

Proof. Let $f(x)=\log(\psi(e^x))$, $x>t=1.1641\ldots$, where t is the solution of the equation $\psi(e^x)=1$. Differentiating with respect to x and by Lemma 2.1 we get

$$\begin{aligned} f''(x) &= \frac{e^x\left(\psi\left(e^x\right)\left(\psi'\left(e^x\right)+e^x\psi''\left(e^x\right)\right)-e^x\psi'\left(e^x\right)^2\right)}{\psi\left(e^x\right)^2}\\ &< \frac{e^x\left(\psi\left(e^x\right)\left(\psi'\left(e^x\right)+1-2e^x\psi'(e^x)\right)-e^x\psi'\left(e^x\right)^2\right)}{\psi\left(e^x\right)^2}\\ &= \frac{e^x\left(\psi\left(e^x\right)(1-(2e^x-1)(1+1/(2e^{2x})))-e^x\psi'\left(e^x\right)^2\right)}{\psi\left(e^x\right)^2}<0. \end{aligned}$$

Hence f is concave, this implies that

$$\frac{\log(\psi(e^x))+\log(\psi(e^y))}{2}\leq\log(\psi(e^{(x+y)/2})).$$

If we let $r=e^x$ and $s=e^y$ we get (1). The proof of part (2) and (3) follow from Lemma 2.8(2). ☐

2.10. **Proof of Theorem 1.4.** Let $f(x)=\log\left(\frac{x\Gamma(x)}{\pi^x}\right)$, $x>1/2$. Differentiating with respect to x and by Lemma 2.1(2) we get

$$\begin{aligned} f''(x) &= \psi'(x)-\frac{1}{x^2} \\ &> \frac{1}{x}+\frac{1}{2x^2}-\frac{1}{x^2}>0. \end{aligned}$$

This implies that the function

$$\log(\Gamma(n/2+1)/\pi^{n/2})=\log(1/\Omega_n),\ n\in\mathbb{N}$$

is convex in n. Hence

$$\log\left(\frac{1}{\Omega_{(x+y)/2}}\right)\le\frac{1}{2}\log\left(\frac{1}{\Omega_x\Omega_y}\right),$$

part (1) follows if we let $x/2=n$ and $y/2=n-1$. Part (2) follows from Lemma 2.5. $\square$

2.11. **Lemma.** [B, Lemma 2.1] *Let us consider the function $f:(a,\infty)\in\mathbb{R}$, where $a\ge 0$. If the function g, defined by*

$$g(x)=\frac{f(x)-1}{x}$$

is increasing on (a,∞), then for the function h, defined by $h(x)=f(x^2)$, we have the following Grünbaum-type inequality

$$1+h(z)\ge h(x)+h(y), \tag{2.12}$$

where $x,y\ge a$ and $z^2=x^2+y^2$. If the function g is decreasing, then inequality (2.12) is reversed.

2.13. **Corollary.** *The following inequalities hold*

(1) $\dfrac{r+s+\psi(r+s)}{r\,\psi(s)+s\,\psi(r)}\ge\dfrac{r+s}{r\,s},\quad r,s\in(c,\infty)$

(2) $\dfrac{1+\Omega_{r+s}}{\Omega_{r+s}}\ge\dfrac{\Omega_r+\Omega_s}{\Omega_r\Omega_s},\quad r,s\in\mathbb{N}\setminus\{1\}.$

Proof. Let

$$f_1(x)=\frac{\psi(x)/x-1}{x},\quad x>c.$$

Differentiating with respect x we get

$$\begin{aligned} f_1'(x) &= \frac{\psi'(x)x+x-2\psi(x)}{x^3} \\ &> \frac{2x^2+2x-4x\log(x)+3}{2x^4}>0, \end{aligned}$$

by Lemma 2.1(1) and (2). Now the proof of part (1) follows from Lemma 2.8(2). For (2), let

$$f_2(x) = \frac{1/\Omega(n) - 1}{n} = \frac{\Gamma\left(n/2\right)}{2\pi^{n/2}} - \frac{1}{n}, \quad x \in \mathbb{N} \setminus \{1\}.$$

By differentiating with respect to n and by Lemma 2.1(1) we get

$$\begin{aligned} f_2'(x) &= \frac{4\pi^{n/2} + n^2\Gamma(n/2)\left(\psi(n/2) - \log(\pi)\right)}{4\pi^{n/2}n^2} \\ &> \frac{4\pi^{n/2} + n^2\Gamma(n/2)\left(\log(n/2) - 2/n - \log(\pi)\right)}{4\pi^{n/2}n^2} > 0. \end{aligned}$$

Hence f_2 is increasing and the proof follows from Lemma 2.8(2). □

References

[AS] M. Abramowitz and I. Stegun, eds.: *Handbook of mathematical functions with formulas, graphs and mathematical tables.* National Bureau of Standards, 1964 (Russian translation, Nauka 1979).

[A1] H. Alzer: *On some inequalities for the gamma and psi functions.* Math. Comp., 66 (1997), 373–389.

[A2] H. Alzer: *Inequalities for the volume of the unit ball in $\mathbb{R}^n$.* J. Math. Anal. Appl. 252 (2000), no. 1, 353–363.

[A3] H. Alzer: *Sharp inequalities for the digamma and polygamma functions*, Forum Math. 16 (2004), 181221

[A4] H. Alzer: *Inequalities for the volume of the unit ball in $\mathbb{R}^n$. II.* Mediterr. J. Math. 5 (2008), no. 4, 395–413.

[AQ] G.D Anderson and S.-L. Qiu: *A monotonicity property of the gamma function*, Proc. Amer. Math. Soc., 125(11) (1997), 3355–3362.

[AVV] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Conformal invariants, inequalities and quasiconformal maps.* J. Wiley, 1997, 505 pp.

[B] Á. Baricz: *Grünbaum-type inequality for general power series.* JIPAM. J. Inequal. Pure Appl. Math. 7 (2006).

[B] N. Batir: *Some new inequalities for gamma and polygamma functions*, J. Inequal. Pure Appl. Math. 6 (2005), no. 4, Art. 103

[EL] Árpád Elbert and Andrea Laforgia : *On some properties of the Gamma function.* Proc. Amer. Math. Soc. 128 (2000), 2667–2673.

[EGP] N. Elezović, C. Giordano and J. Pečarić: *The best bounds in Gautschis inequality,* Math. Inequal. Appl. 3 (2000), 239252.

[KMSV] R. Klén, V. Manojlović, S. Simić and M. Vuorinen: *Bernoulli inequality and hypergeometric functions.* Proc. Amer. Math. Soc. (to appear) arXiv math.CA 1106.1768.

[M] C. Mortici: *Monotonicity properties of the volume of the unit ball in $\mathbb{R}^n$.* (English summary) Optim. Lett. 4 (2010), no. 3, 457–464.

[N] E. Neuman: *Inequalities involving a logarithmically convex function and their applications to special functions.* J. Inequal. Pure Appl. Math. Article 16, 2006.

[QV] S.-L. Qiu and M. Vuorinen: *Some properties of the gamma and psi functions, with applications.* Math. Comp. 74 (2005), no. 250, 723–742.

[R] H. Ruskeepää: *Mathematica® Navigator.* 3rd ed. Academic Press, 2009.

[S] J. Sándor: *A note on certain inequalities for the gamma function.* J. Ineq. Pure Appl. Math., 6(3) (2005), Art. 61.

E-mail address: `bhayo.barkat@gmail.com`